

Quantum friction on atoms after acceleration

Diego A. R. Dalvit
Theoretical Division
Los Alamos National Laboratory


Work done in collaboration with
F. Intravaia, R. Behunin, V. Mkrtchian, S. Buhmann, S. Schell, C. Henkel

PRA 89, 050101(R) (2014)

arXiv:1502.01117, to appear in JPCM




Outline of this talk




- What is quantum friction?
- Quantum friction from non-equilibrium FDT
- Quantum friction from t-dep. perturbation theory
- Key results
 - Quantum friction depends on the way the atom is boosted
 - Quantum friction is cubic in velocity

An intuitive picture




An intuitive picture



An intuitive picture


Photons and plasmon field
perceived with a Doppler
shifted frequency



A variety of predictions

Zero temperature atom-surface quantum friction

Authors	Low velocity dependency	Distance dependency	Comments
Mahanty 1980	\mathbf{v}	\mathbf{z}^{-5}	Approach similar to the calculations of vdW forces but with mistakes
Schaich and Harris 1981	\mathbf{v}	\mathbf{z}^{-10}	Two-state atom with a transition dipole moment normal to a metal surface
Scheel and Buhmann 2009	\mathbf{v}	\mathbf{z}^{-8}	Master-equation approach for multilevel atoms and quantum regression “theorem”.
Barton 2010	\mathbf{v}	\mathbf{z}^{-8}	Perturbation theory using Fermi’s golden rule. Harmonic oscillator.
Philbin and Leonhardt 2009	$\mathbf{0}$	$\mathbf{0}$	Relativistic calculations and analytical/numerical evaluation of the Green’s tensor
Dedkov and Kyasov 2012	\mathbf{v}^3	\mathbf{z}^{-7}	Fluctuation-dissipation theorem applied to the dipole atom as well as to the electric field

$$F_{\text{fric}}(t) = \langle \hat{\mathbf{d}}(t) \cdot \partial_x \hat{\mathbf{E}}(\mathbf{r}_a(t), t) \rangle \quad \longleftrightarrow \quad F_{\text{ext}}(t)$$


Prescribed motion

$$\mathbf{r}_a(t) = \begin{cases} (x_0, y_a, z_a) & \text{for } t < t_a \\ (x_{\text{accel}}(t), y_a, z_a) & \text{for } t_a < t < 0 \\ (x_a + v_x t, y_a, z_a) & \text{for } t > 0 \end{cases}$$

$$m_a \ddot{x}_a(t) = F_{\text{ext}}(t) + F_x(t)$$

Stationary ($t \rightarrow \infty$) frictional force

$$F_{\text{fric}} = \text{Re} \left\{ \frac{2}{\pi} \int \frac{d^2 \mathbf{k}}{(2\pi)^2} k_x \int_0^\infty d\omega \int_0^\infty d\tau e^{-i(\omega - k_x v_x)\tau} \text{Tr}[\underline{C}(\tau; v_x) \cdot \underline{G}_I(\mathbf{k}, z_a, \omega)] \right\}$$

$$\underline{C}_{ij}(\tau; v_x) = \text{tr} \{ \hat{d}_i(0) \hat{d}_j(-\tau) \hat{\rho}(\infty) \}$$

No general results for non-equilibrium state $\hat{\rho}(\infty) = ???$

Non-equilibrium FDT

● Dipole moment $\hat{\mathbf{d}} = \mathbf{d}\hat{q}$ $\ddot{\hat{q}}(t) + \omega_a^2 \hat{q}(t) = \frac{2\omega_a}{\hbar} \mathbf{d} \cdot \hat{\mathbf{E}}(\mathbf{r}_a(t), t)$

● Dynamic polarizability of moving atom

$$\underline{\alpha}_{ij}(\omega; v_x) = \frac{2\omega_a}{\hbar} \mathbf{d}_i \mathbf{d}_j \left[-\omega^2 + \omega_a^2 - \frac{2\omega_a}{\hbar} \int \frac{d^2 \mathbf{k}}{(2\pi)^2} \mathbf{d} \cdot \underline{G}(\mathbf{k}, \omega + k_x v_x) \cdot \mathbf{d} \right]^{-1}$$

● An exact, non-equilibrium fluctuation-dissipation relation

$$\underline{S}(\omega; v_x) = \frac{\hbar}{\pi} \theta(\omega) \underline{\alpha}_I(\omega; v_x) - \frac{\hbar}{\pi} \underline{J}(\omega; v_x)$$

Non-equilibrium FDT in classical models have the same form

(Chetrite et al. 2008)

$$\underline{J}(\omega; v_x) = \int \frac{d^2 \mathbf{k}}{(2\pi)^2} [\theta(\omega) - \theta(\omega + k_x v_x)] \underline{\alpha}(\omega; v_x) \cdot \underline{G}_I(\mathbf{k}, \omega + k_x v_x) \cdot \underline{\alpha}^*(\omega; v_x).$$


● Using $\underline{S}(\omega; v_x)$ one can obtains

$$F_{\text{fric}} \approx -\frac{45\hbar}{256\pi^2 \epsilon_0} \alpha'_I(z_a, 0) \Delta'_I(0) \frac{v_x^3}{z_a^7}$$

Other approach: pert. theory

- Multi-level atom, initially in its ground state
- Polariton field (EM+matter) also in its ground state

$$\rho(0) = |g\rangle\langle g| \otimes |\text{vac}\rangle\langle \text{vac}|$$




- Prescribed motion $\mathbf{r}_a(t) = \begin{cases} (x_0, y_a, z_a) & \text{for } t < t_a \\ (x_{\text{accel}}(t), y_a, z_a) & \text{for } t_a < t < 0 \\ (x_a + v_x t, y_a, z_a) & \text{for } t > 0 \end{cases}$

- Our goal is to calculate:

radiative frictional force \mathbf{F}

radiative frictional power $P = -\mathbf{v} \cdot \mathbf{F}$



Transition amplitudes

- Polariton field Hamiltonian (near-field regime)

$$\hat{\Phi}(\vec{r}, t) = \int d^2k \int_0^\infty d\omega (\hat{a}_{\mathbf{k}\omega} \phi_{\mathbf{k}\omega} \exp(i\mathbf{k} \cdot \mathbf{r} - i\omega t) + \text{h.c.}) \quad \phi_{\mathbf{k}\omega} = \frac{\sqrt{\omega \Gamma \omega_p^2/2}}{\omega^2 + i\omega \Gamma - \omega_S^2} \sqrt{\frac{\hbar}{2\pi^2 k}} e^{-kz}$$


- Atom-field coupling $\hat{V}(t) = -\hat{D}_i(t) \hat{E}_i(\vec{r}(t), t) = \hat{D}_i \partial_i \hat{\Phi}(\vec{r}(t), t)$

- Atom-field states $|\Psi(t)\rangle = (1 + c_0^{(2)}(t))|g, \text{vac}\rangle + \sum_{\vec{\eta}} \int d^3\kappa (c_1^{(1)}(t) + c_1^{(3)}(t))|\vec{\eta}, \kappa\rangle$

(third order expansion in powers of d)

$$c_n^{(p)}(t)$$

transition amplitudes for states with n-photons in the p-th pert. order



$$\begin{aligned} \langle g, \text{vac} | \hat{V}(t) | \vec{\eta}, \kappa \rangle &= i d (\vec{\eta} \cdot \vec{k}) \phi_{\kappa} \exp[-i(\Omega + \omega)t + i\mathbf{k} \cdot \mathbf{r}(t)] , \\ \langle \vec{\eta}, \kappa | \hat{V}(t) | g, \kappa_1 \kappa_2 \rangle &= i d (\vec{\eta} \cdot \vec{k}_1) \phi_{\kappa_1} e^{i(\Omega - \omega_1)t + i\mathbf{k}_1 \cdot \mathbf{r}(t)} \delta(\kappa - \kappa_2) \\ &\quad + i d (\vec{\eta} \cdot \vec{k}_2) \phi_{\kappa_2} e^{i(\Omega - \omega_2)t + i\mathbf{k}_2 \cdot \mathbf{r}(t)} \delta(\kappa - \kappa_1) , \\ \langle \vec{\eta}, \text{vac} | \hat{V}(t) | g, \kappa \rangle &= i d (\vec{\eta} \cdot \vec{k}) \phi_{\kappa} \exp[i(\Omega - \omega)t + i\mathbf{k} \cdot \mathbf{r}(t)] . \end{aligned}$$

Frictional force and power

Quantum friction force operator:

$$\hat{\mathbf{F}}(t) = \int d^2k \int_0^\infty d\omega \left(\mathbf{k}(\vec{\tilde{D}}(t) \cdot \vec{k}) \phi_{\mathbf{k}\omega} \hat{a}_{\mathbf{k}\omega} \exp(i\mathbf{k} \cdot \mathbf{r}(t) - i\omega t) + \text{h.c.} \right)$$

Radiated power:

$$P = P_1 + P_2$$

$$P_1 = \lim_{t \rightarrow \infty} \sum_{\vec{\eta}} \int d^3\kappa \hbar(\Omega + \omega) \frac{|\langle \vec{\eta}, \kappa | \Psi(t) \rangle|^2}{t}$$

$$P_2 = \frac{1}{2} \lim_{t \rightarrow \infty} \int d^3\kappa_1 \int d^3\kappa_2 \hbar(\omega_1 + \omega_2) \frac{|\langle g, \kappa_1 \kappa_2 | \Psi(t) \rangle|^2}{t}$$

- To second order in perturbation theory, the force and power are exponentially small

$$\mathbf{F}^{(2)} \propto \exp(-2\Omega z/v_x)$$

$$P^{(2)} \propto \exp(-2\Omega z/v_x)$$

- To fourth order in perturbation theory, great care must be exercised because of possible interferences between the amplitudes $c_n^{(p)}(t)$

Ideal case: constant v always

We consider first the idealized case of the atom moving at constant velocity at all times $\mathbf{r}(t) = \mathbf{v}t$, $\forall t$

One-photon power dissipated P_1

$$P_1^{(4)} \rightarrow 2 \operatorname{Re}[c_1^{(1)*}(t)c_1^{(3)}(t)] \approx -\gamma_g t P_1^{(2)}$$

hence, it is exponentially suppressed

Two-photon power dissipated P_2

$$P_2^{(4)} \rightarrow |c_2^{(2)}|^2 \propto \frac{v_x^4}{z_a^{10}}$$


Frictional force

$$F_x^{(4)} \propto \frac{v_x^3}{z_a^{10}}$$

Boosting the atom: $v(t)$

We now consider the more realistic case in which the atom is boosted from its initial rest position to a final state in which it moves at constant velocity

Curve	$\dot{x}(t)$	$\ddot{x}(t)$
thick black	$v/(1 + e^{-t/\tau})$	$v\tau^{-1}/(2 + 2 \cosh t/\tau)$
thin blue	$\begin{cases} 0 & \text{for } t < -\tau \\ (t + \tau)v/(2\tau) & \text{for } -\tau < t < \tau \\ v & \text{for } t > \tau \end{cases}$	$\begin{cases} 0 & \text{for } t < -\tau \\ v/(2\tau) & \text{for } -\tau < t < \tau \\ 0 & \text{for } t > \tau \end{cases}$
dashed gray	$\begin{cases} 0 & \text{for } t < 0 \\ v & \text{for } t > 0 \end{cases}$	$v\delta(t)$



Two-photon power $P_2^{(4)} = P_A + P_B$
 $P_A \propto v_x^4$ is independent of the boost \leftarrow (Barton 2010)

$P_B \propto v_x^2 f(\tau)$ depends on the boost, and is exponentially suppressed for adiabatic boosts $\omega\tau \gg 1$

One-photon power $P_1^{(4)} = -P_B \leftarrow$ subtle cancellation!

Hence, perturbation theory also gives cubic-in- v quantum friction

Orders of magnitude

● Near-field quantum friction

$$F_{\text{fric}} \approx - \frac{45\hbar\rho^2\alpha_0^2}{512\pi^3} \frac{v_x^3}{z_a^{10}}$$

surface's electrical resistivity static atomic polarizability

● Example: ground state ^{87}Rb flying over a silicon surface

$$\alpha_0 = 5.26 \times 10^{-39} \text{ Hz}/(\text{V}/\text{m})^2$$

$$\rho = 6.4 \times 10^2 \text{ } \Omega \text{ m}$$

$$v_x = 340 \text{ m/s}$$

$$z_a = 10 \text{ nm}$$

$$\Rightarrow F_{\text{fric}} \approx -1.3 \times 10^{-20} \text{ N}$$



● How to enhance it? How to measure it?

- excited atomic states?
- higher velocities?
- materials with higher resistivities?
- macroscopic bodies?
- ???
- atomic interferometry?
- near-field AFM?
- ???

Conclusions

- Atom-surface quantum friction from general non-equilibrium stat. mech.
- Non-equilibrium FDT predicts a cubic-in- v frictional force
- Atom-surface quantum friction from t -dependent perturbation theory
 - Dependency on boost history
 - Subtle cancellation between one- and two-photon dissipated power. This results, again, in cubic-in- v quantum friction
- At high temperatures (classical limit), linear-in- v frictional force
- Same analysis possible for quantum friction between macroscopic bodies